

Digital Signal Processing Techniques: Calculating Biological Functionalities

Norbert Nwankwo* and Huseyin Seker

The Bio-Health Unit of the Centre for Computational Intelligence, De Montfort University, Leicester, United Kingdom

Abstract

Calculating Biological functionalities of proteins and presenting them numerically **is an** approach that will benefit the designing of drugs and vaccines. For example, potency of vaccines is known to be measured in terms of specificity, which is determined by bio-recognition (affinity), and sensitivity. Calculating the bio-recognition of peptides employed in the design of vaccines by means of procedures like the Digital Signal Processing (DSP) techniques rather than clinical experimentations not only gives room for the manipulation of the amino acids sequences of the peptides but also helps determine the degree of specificity of the antibody generated, hence the potency of the vaccine produced. This also provides opportunity for optimization of the peptides for the desired biological characteristics. Also comparing the potency of peptide-based drugs through calculation of their biological functions is a faster, easier and resource-saving approach to pharmacotherapeutic investigations.

In this study, two DSP techniques are fully explained and demonstrated. They are Resonant Recognition Model and Informational Spectrum Method. They are employed in the calculation of some physiological characteristics of Plasmodial peptides (P18 and P32), which are still being study for possible use as materials for the designing of anti-Malaria vaccines. Furthermore, the approaches are utilised to assess the pharmacological activities of two Fusion inhibitors (Enfuvirtide and Sifuvirtide). Enfuvirtide is currently in use for the management of anti-HIV/AIDS while Sifuvirtide is still being studied.

Our calculated results demonstrate strong correlation with the preliminary clinical findings. They also seem to suggest that presenting biological characteristics in numerical terms is an easier and more rational approach to designing drugs and vaccines as it save resources and time unlike clinical experimentations. The methods also appear to help simplify the manoeuvring of the protein residues, which are employed in the designing and development of drugs and vaccines in order to obtain maximal biological characteristics.

Keywords: Digital signal processing; Fusion peptide; Informational spectrum method; Resonant recognition model

Introduction

Digital Signal Processing techniques are analytic procedures, which decompose and process signals in order to reveal information embedded in them [1]. The signals may be continuous (unending) or discrete such as the protein residues. Digital Signal Processing techniques have helped analyse protein interactions [2] and made biological functionalities calculable including drug resistance [3]. This is in contrast to Bioinformatics methods like Multiple Structural Alignment (MSA), which dwells on homology and as such predict protein functionalities.

In these approaches, protein residues are first converted into numerical sequences (signals) using one out of over 600 available amino acids parameters that are responsible for the biological functionality. These numerical sequences (signals) are then processed by means of Discrete Fourier Transform (DFT) to present the biological characteristics of the proteins in the form of Informational Spectrum. This procedure is called Informational Spectrum Method (ISM) [4-6].

A version of the ISM, which engages amino acids parameter called Electron-Ion Interaction Potential (EIIP) is referred Resonant Recognition Model (RRM). In this procedure, biological functionalities are presented as Spectral Characteristics [2]. This physico-mathematical process, which is based on the fact that bio-molecules with same biological characteristics recognise and bio-attach to themselves when their valence electrons oscillate and then reverberate in an electromagnetic field [2].

In this study, detailed explanation of Resonant Recognition Model and Informational Spectrum Method (ISM) are provided. Furthermore, demonstration of these steps is made using one out of

over 600 amino acids parameters, and two Plasmodial peptides (P18 and P32) [4]. Thereafter, the techniques are used to calculate biological functionalities of peptide-based anti-HIV /AIDS drugs (Enfuvirtide and Sifuvirtide) and the two Plasmodial peptides (P18 and P32) which are presented numerically in order to study their functionalities.

Resonant Recognition Model (RRM): Background

Resonant Recognition Model (RRM) is a Digital Signal Processing-based technique, which recognizes protein primary structures or physiological functionalities as protein residues represented by numbers that are assigned from the Electron-ion Interaction Pseudopotential (EIIP) parameter [2]. RRM involves four steps, which are fully discussed below.

The steps include:

Step 1: Conversion of the Protein Residues into Numerical Values Electron-ion Interaction Pseudopotential (EIIP) Parameter: There are 20 essential amino acids constituents (protein residues) [7]. Biological interactions have been studied in relation to the behaviour of these 20 amino acids constituent of proteins. As a

*Corresponding author: Norbert Nwankwo, The Bio-Health Unit of the Centre for Computational Intelligence, De Montfort University, Leicester, United Kingdom, E-mail: nnwankwo@dmu.ac.uk or nnwankwo@hotmail.co.uk

Received September 20, 2011; Accepted November 03, 2011; Published December 01, 2011

Citation: Nwankwo N (2011) Digital Signal Processing Techniques: Calculating Biological Functionalities. J Proteomics Bioinform 4: 260-268. doi:10.4172/jpb.1000199

Copyright: © 2011 Nwankwo N. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

result, the level of participation by these protein residues in over 565 protein-protein interactions that characterise biological functionalities have been derived as Amino Acids Parameters (AAPs) and deposited in databases like www.genome.jp/aaindex/ [8] and literatures such as [9]. AAPs describe physiochemical characteristics such as Hydropathy [10]; Hydrophobicity [11,12]; Hydrophilicity [13,14]; and Structural features including Alpha-Helix [15,16], and Beta [16] conformations.

Electron-Ion Interaction Potential (EIIP) (Table 1) is one of the amino acid parameter. The Resonant Recognition Model (RRM) engages EIIP amino acids parameter. Proteins molecules are known to be sequences of amino acids with unrestricted electrons and charges [17]. These charges elicit short-lived polarization of the side-chain groups and result in electromagnetic oscillation between some parts of the protein molecule [18]. These swinging in the protein molecules interfere with one another [19,20]. During oscillation, molecules, which share same biological characteristics, are found to resonate at the same frequency leading to amplified attraction (affinity) [18-20]. This reverberation, which arises from electromagnetic oscillation between bio-molecules (electromagnetic resonance) [2], is called the Electron-Ion Interaction Potential (EIIP) [19,21]. RRM which engages EIIP and therefore determines biological characteristics of the protein residues in terms of bio-recognition (specificity) and binding interaction (affinity) [2]. Bio-recognition and bio-attachment are the first two steps in molecular interactions. By means of DSP, the biological functionalities of the peptide-based anti-HIV/AIDS drugs and Plasmodial peptides are calculated. This is in an attempt to find out if this rational approach is realizable and beneficial. Plasmodial peptides are prospective starter materials for Malaria vaccine design peptides.

Step 2: Zero-padding/Upsampling: In some cases, proteins that are to be analyzed by means of RRM may have unequal residues. Signal Processing techniques demand that the window length of all proteins be the same [1,28]. Such unequal window lengths can be found in Plasmodial peptides P18 and P32 being used in this study as shown in Table 2. P18 has 16 amino acids compositions while P32 has the largest protein length (N), which is 32. Therefore, prior to the

Amino Acid	Three Letter Code	Letter Code	EIIP
Alanine	Ala	A	0.0373
Arginine	Arg	R	0.0959
Asparagine	Asn	N	0.1263
Aspartic acid	Asp	D	0.0036
Cysteine	Cys	C	0.0829
Glutamine	Gln	Q	0.0761
Glutamic acid	Glu	E	0.0058
Glycine	Gly	G	0.0050
Histidine	His	H	0.0242
Isoleucine	Iso	I	0.0000
Leucine	Leu	L	0.0000
Lysine	Lys	K	0.0371
Methionine	Met	M	0.0823
Phenylalanine	Phe	F	0.0946
Proline	Pro	P	0.0198
Serine	Ser	S	0.0829
Threonine	Thr	T	0.0941
Tryptophan	Trp	W	0.0548
Tyrosine	Tyr	Y	0.0516
Valine	Val	V	0.0057

Table 1: Amino Acids Parameter: Electron-ion Interaction Pseudopotential (EIIP)

decomposition of the numerical signals by means Fourier Transform (FT), the sequences are to be brought to same sequence (window) length with largest protein residue (N). In the case of peptides P18 and P32, 14 zeros are added to peptide P18. This is called zero-padding or up-sampling [22]. Zero-padding operation is usually used to improve the visual clarity of the spectrum. It does not improve the quality of the spectral results. However, zero-padding has been claimed to constitute error [23].

Step 3: Processing of the Numerical Sequences (Signals) using Fourier Transform (FFT): Fourier Transform is a mathematical operation that converts one signal to another without altering the information contained in the signal [1]. This is in an attempt to disclose hidden information. Because proteins are amino acids in compartments (digitalized), Discrete Fourier Transform (DFT) is employed in their analysis [1]. Though DFT is known to be the most straight-forward mathematical procedure [1], it is found to be inefficient.

As a result, a more efficient, rational and faster algorithm, which executes DFT faster without altering the results, is being employed [24]. It is called Fast Fourier Transform (FFT), and is represented as:

$$y = \text{abs}(\text{fft}(x)) \tag{1}$$

RRM-based decomposed signals results in three sets of values namely imaginary, real and absolute. The plots of these values describe the Spectral Characteristics (SC) of proteins. The Absolute values have been exploited in assessing biological functionalities of proteins. However, Imaginary and Real values have recently been found to reveal some biological characteristics [25]. The y-axis (Amplitude) symbolizes the bio-recognition (specificity) and binding (affinity) of each protein residue while the x-axis (Frequency) determines the position of the interaction.

Step 4: Cross-spectral Analysis: In order to identify the common biological relationship amongst proteins, point-wise multiplication of the Spectral Characteristics of the protein residues are applied [26]. This process is called Cross-spectral Analysis [2]. Proteins with common biological functionality are known to share one significant peak, called the Consensus Frequency [2], which is acknowledged to represent the region responsible for the biological functionality.

Analysis of the Results: Analysis of the result entails identifying the Consensus Frequency (CF) and the utilisation of the amplitude of the Spectral Characteristics of the protein residues at the CF to study their physiological properties. According to the RRM procedure [2], the relationship between the Consensus Frequency (F) and the Peak Position (PP) can be expressed as:

$$CF = \frac{PP * 0.5}{N / 2} \tag{2}$$

This is same as:

$$CF = \frac{PP}{N} \tag{3}$$

where N represents the length of the largest protein in the dataset.

The amplitudes attained by protein residues at the CF demonstrated are useful in determining relationships that exist between the proteins as well as the organisms that harbour them [5,27]. Results can be expressed in terms of percentages depending on the scaling factor. When the scaling function is the maximum value, the highest amplitude obtainable is 1 or 100%.

Methodology

Method 1: Demonstration of Resonant Recognition Model

Plasmodial peptides P18 and P32 shown in Table 2 have clinically been found to provide immunization against *Plasmodium berghei* in rodents [28]. They are unequal in length. While peptide P18 has 18 amino acids components, peptide 32 has 32 residues. Using EIIP parameter that is concerned with bio-recognition (specificity of the antibody produced by the peptides) and bio-attachment (relating to the sensitivity of the antibody), RRM procedure is first demonstrated and then used to calculate the degree of bio-recognition and binding interaction. Potency of a vaccine is determined by its specificity and

Peptide Identity	Peptide
P18	EWSPCSVTCGNGIQVRIK
P32	IEQYLKKIKNSISTEWSVTCGNGIQVRIK

Table 2: Peptides for the Demonstration of RRM and ISM.

S/No	Seq (P32)	EIIP	CS	CF	Seq (P18)	EIIP	SC	CF
1	I	0.0000	-		E	0.0058	-	-
2	E	0.0058	0.6325		W	0.0548	0.4778	0.3022
3	Q	0.0761	0.4019		S	0.0829	0.6211	0.2496
4	Y	0.0516	0.1830		P	0.0198	0.5836	0.1068
5	L	0.0000	0.6461		C	0.0829	0.4260	0.2752
6	K	0.0371	0.3963		S	0.0829	0.5025	0.1992
7	K	0.0371	0.3891		V	0.0057	0.6171	0.2401
8	I	0.0000	0.2667		T	0.0941	0.1758	0.0469
9	K	0.0371	0.4000		C	0.0829	0.0920	0.0368
10	N	0.1263	0.2810		G	0.0050	0.5965	0.1676
11	S	0.0829	0.7978		N	0.1263	0.6975	0.5564
12	I	0.0000	0.5506		G	0.0050	0.7390	0.4069
13	S	0.0829	1.0000		I	0.0000	1.0000	1.0000
14	T	0.0941	0.3502		Q	0.0761	0.4411	0.1545
15	E	0.0058	0.3770		V	0.0057	0.5383	0.2029
16	W	0.0548	0.7558		R	0.0959	0.5850	0.4422

Table 3: Summary of the RRM Procedure: P16 and P32.

S/No	Seq (P32)	EIIP	CS	CF	Seq (P18)	EIIP	SC	CF
17	S	0.0829	0.7558		I	0.0000	0.5850	0.4422
18	P	0.0198	0.3770		K	0.0371	0.5383	0.2029
19	C	0.0829	0.3502		-	0	0.4411	0.1545
20	S	0.0829	1.0000		-	0	1.0000	1.0000
21	V	0.0057	0.5506		-	0	0.7390	0.4069
22	T	0.0941	0.7978		-	0	0.6975	0.5564
23	C	0.0829	0.2810		-	0	0.5965	0.1676
24	G	0.0050	0.4000		-	0	0.0920	0.0368
25	N	0.1263	0.2667		-	0	0.1758	0.0469
26	G	0.0050	0.3891		-	0	0.6171	0.2401
27	I	0.0000	0.3963		-	0	0.5025	0.1992
28	Q	0.0761	0.6461		-	0	0.4260	0.2752
29	V	0.0057	0.1830		-	0	0.5836	0.1068
30	R	0.0959	0.4019		-	0	0.6211	0.2496
31	I	0.0000	0.6325		-	0	0.4778	0.3022
32	K	0.0371	-		-	0	-	-

Table 4: Summary of the RRM Procedure: P16 and P32 (continue).

Step 2: Conversion of Alphabetic codes into Numerical Signals using EIIP

> Numerical Sequence of P18.

0.0058, 0.0548, 0.0829, 0.0198, 0.0829, 0.0829, 0.0057, 0.0941, 0.0829, 0.0050, 0.1263, 0.0050, 0.0000, 0.0761, 0.0057, 0.0959, 0.0000, 0.0371.

> Numerical of Sequence of P32

0.0000, 0.0058, 0.0761, 0.0516, 0.0000, 0.0371, 0.0371, 0.0000, 0.0371, 0.1263, 0.0829, 0.0000, 0.0829, 0.0941, 0.0058, 0.0548, 0.0829, 0.0198, 0.0829, 0.0829, 0.0057, 0.0941, 0.0829, 0.0050, 0.1263, 0.0050, 0.0000, 0.0761, 0.0057, 0.0959, 0.0000, 0.0371

Figure 1: EIIP-based Numerical Sequences of P18 and P32.




Figure 2: Plot of EIIP-based Numerical Sequence of P18 and P32 (without padding).

sensitivity. In this study, Plasmodial peptides P18 and P32 are used to demonstrate the RRM procedures.

Demonstration Step 1: Conversion of the Protein Residues into Numerical Values of EIIP Parameter: The entire RRM process is demonstrated in Tables 3 and 4.

The Alphabetic codes in the sequences of the peptides P18 and P32 demonstrated in Table 2 are interchanged with the corresponding EIIP values shown in Table 1 in order to obtain their numerical sequences

to 32 of the P18 and as such, they are zero-padded. The sequences are then translated using EIIP, processed by means DFT to yield Spectral Characteristics (SC) and point-wise multiplied to generate the Cross Spectral (CS) features.

As demonstrated in Table 3, the point-wise multiplication at position 2 of the Spectral Characteristics of P32 (0.6325) and P18 (0.4778) yields 0.3022. This process is carried out all through the sequences. The plot of the CS values yields a symmetric image Figure 4 which is halved and scaled with 0.5. Also, "DC component" of the CS is discarded. Therefore, out of the 32 sequences, 15 values of the Cross Spectral (CS) features are also obtained and illustrated in Figure 5. The plot of the Spectral Characteristics of P18 and P32 are shown in Figure 6 while their Cross-Spectral Characteristics is displayed in Figure 7

Consensus Frequency: According to the RRM procedure [2], the relationship between the Consensus Frequency (F) and the Peak Position (PP) can be expressed as:

$$CF = \frac{PP * 0.5}{N / 2} \quad (5)$$

This is same as:

$$CF = \frac{PP}{N} \quad (6)$$

where N represents the length of the largest protein in the dataset.

$$CF = \frac{12}{32} \quad (7)$$

Based on equation 7, the CF of P18 and P32 is 0.375.

The Consensus Frequency for the two the Plasmodial peptides P18 and P32 is 0.375 (Position 12).

Using the amplitude value of the proteins and peptides at the CF, biological relationships between the proteins analysed can be identified and their biological functionalities calculated.

Method 2: Information Spectrum Method (ISM)

Information Spectrum Method (ISM), like the RRM is a Digital Signal Processing-based technique that considers protein primary structures or physiological functionalities amino acids sequences of protein represented by numbers using, unlike the RRM, any of the Amino Acids Parameter [2,6,30]. Graphic representation of the biological characteristics by means ISM is referred to as




Figure 7: Cross-Spectral Spectral Characteristics of P18 and P32.




Figure 8: CHAM830107-based Information Spectra (IS) of P18 and P32.

the Informational Spectrum (IS) [2]. Furthermore, point-wise multiplication of the Spectral Characteristics results which reveals shared biological functionalities by proteins demonstrated as a prominent peak called Consensus Frequency (CF) is referred to as Common Informational Spectrum (CIS) [2]. ISM procedure has been used to investigate principal arrangement in Calcium binding protein [30], Influenza viruses [6].

Information Spectrum Method (ISM) uses the same procedure as the RRM. It involves three main steps. Like the RRM, the steps include the conversion the alphabetic code of amino acids sequences into numerical values using amino acids scale that relates to the interaction under investigation. This is followed by the processing of the numerical sequences (signals) using discrete Fourier Transform (DFT). Absolute values of the complex DFT represented as a plot called Informational Spectrum (IS) discloses the information embedded in the protein residues. The y-axis (Amplitude) signifies the contribution in terms of susceptibility or resistance by each sequence while the x-axis (Frequency) determines the position of biological interaction. The third step entails obtaining Common Informational Spectrum (CIS). CIS compares the activities of proteins, which have common biological functions. CIS is the point-wise multiplication of Informational Spectrum (IS).

Demonstration of Informational Spectrum: ISM and the RRM procedure are same. Like in the case of RRM, ISM procedure starts with the conversion of the alphabetic codes of the P18 and P32 peptide residues as shown in Figure 2 into numerical sequences using CHAM830107 parameter in Table 5. The CHAM830107 translated

sequences obtained thereafter are then processed by means of Discrete Fourier Transform to obtain Informational Spectra (IS) of P18 and P32.

Using equation 6, the Consensus Frequency is recognized as:

$$CF = \frac{3}{32} = 0.094 \quad (8)$$

Therefore, the CF of the two Plasmodial peptides based on CHAM830107 is 0.094.

Based on the amplitudes of the protein residues at the CF identified, further investigations into the biological behaviour of the proteins can be disclosed by means of the ISM and amino acids parameter, CHAM830107 engaged.

Materials

Amino acids sequences of the Plasmodial Circumsporozoite, HIV glycoprotein41 from HXB2 isolate are retrieved from UNIPROT [31]. P18, P32, and Enfuvirtide and Sifurvide are obtained from the literature. While P18 and P32 amino acids sequences are recognized as being studied as starter materials for vaccine development [28], the Enfuvirtide amino acids sequence has been examined along with that of the Sifuvirtide, which is noted to be a bio-medically engineered peptide, obtained from the HIV gp41 C-terminal of the Heptad Repeat (CHR) sequence of the HIV subtype E [13,32].

Results and Discussions

Preliminary Clinical Studies: P18 and P32

The two sets of pharmacologically active components are the Plasmodial peptides P18 and P32, as well as anti-HIV Fusion peptides namely; Enfuvirtide (T20) and Sifuvirtide are first clinically studied. Using the related Amino Acids Parameters, these pharmacologically activities are calculated by means of Digital Signal Processing technique and correlated with our computational outcomes.

Peptides P18 and P32 from the Plasmodium *falciparum* have been identified to inhibit Plasmodium *berghei* invasion of Hep-G2 while P32 is found to protect immunised mice [28]. Interactions between the CS and the hepatocytes and subsequent invasion of the Hep-G2

by the Plasmodial sporozoites have been recognised [33,34]. Prior to interaction, the two proteins must bio-recognise and bind. This interaction is governed by EIIP parameter [35]. Negatively charged carboxyl group of the GAGs have been found to partake in the binding of the CS to HSGP [26]. As a result, amino acids parameters that engage charges (Positive and Negative) are employed in this study.

Results and Discussions: P18 and P32

In this study, EIIP (Table 2) and five (5) other amino acids parameters such as the amino acids parameter with descriptor CHAM830107 (Table 5), which are based on charges including CHAM830108, FAUJ880111, FAUJ880112 and KLEP840101 are retrieved from [16] and engaged in the demonstration of the ISM analysis of the P18 and P32 peptides sequences shown in Table 2 and further used to calculate their biological functionalities. The results are presented in Table 6.

As shown in Figures 6 and 7, both P18 and P32 have maximum amplitudes of 1.00 at the CF, which is at Position 12. This appears to indicate 100% bio-recognition and bio-attachment for the two peptides (P18 and P32). This characteristic determines specificity of interaction between the peptides and the ligands that produce the immunizing antibodies, a property that determines vaccine potency.

It is therefore demonstrated in this study that both P18 and P32 have amplitude of 1.00, which suggest 100% affinity for the target protein that will elicit the production of the antibody necessary for the immunization against Malaria. These are peptides that are still being studied for possible engagement in the designing of vaccines. Vaccines are required to demonstrate specificity in action so that they could produce specific antibody that would neutralise specific antigen. The outcome of this study appears to suggest that both peptides are suitable starter materials for the design of Malaria vaccines by their specificity (precision in the production of Malaria vaccine rather than any other disease) as demonstrated in their ability to bio-identify and bio-attach to the target protein.

Though the results of the EIIP (Figure 6) and FAUJ880111 (Table 6) parameters demonstrated equi-potency by maintaining same maximum (100%) interaction (Figure 2 and Table 6), other parameters disclose unequal potencies. Using CHAM830107 and CHAM830108, our results reveal higher and maximum (100%) interaction for the P18 (Table 6). For the P32, the CHAM830107 parameter yields 89.8% interaction while the CHAM830108 produces 81.1% interaction.

On the other hand, the FAUJ880112 and KLEP840101-based analyses show higher and maximum (100%) for the P32. These seem to indicate higher sensitivity; hence potency for the P32 peptides in terms of the FAUJ880112 and KLEP840101 parameters. As shown in Table 6, FAUJ880112 parameter revealed 100% interaction for the P32 at the CF and 71.6% for the P18. In the same manner, KLEP840101-based analysis reveals 100% interaction for the P32 and 88.8% for the P18. These appear to suggest different sensitivity and as such, different capability for the antibody generated in immunizing host organisms for Malaria except for the EIIP and FAUJ880111 which demonstrated equi-potency.

The six amino acids parameters studied and shown in Table 6 therefore express the specificity (bio-recognition and bio-attachment to the target proteins that direct the production of antibody for Malaria only). In the case of EIIP parameter which measures specificity, the level of protection from the Malaria (potency) by the two Plasmodial peptides used (P18 and P32) is the same while the degree of sensitivity as provided by the other parameters vary. These physiologic indices

Amino Acid	Three Letter Code	One Letter Code	CHAM830107
Alanine	Ala	A	0.00
Arginine	Arg	R	0.00
Asparagine	Asn	N	1.00
Aspartic acid	Asp	D	1.00
Cysteine	Cys	C	0.00
Glutamine	Gln	Q	0.00
Glutamic acid	Glu	E	1.00
Glycine	Gly	G	1.00
Histidine	His	H	0.00
Isoleucine	Iso	I	0.00
Leucine	Leu	L	0.00
Lysine	Lys	K	0.00
Methionine	Met	M	0.00
Phenylalanine	Phe	F	0.00
Proline	Pro	P	0.00
Serine	Ser	S	0.00
Threonine	Thr	T	0.00
Tryptophan	Trp	W	0.00
Tyrosine	Tyr	Y	0.00
Valine	Val	V	0.00

Table 5: A parameter of charge transfer capability (CHAM830107).

Sequence	EIIP	CHAM830107	CHAM830108	FAUJ880111	FAUJ880112	KLEP840101
P16	1.000	1.000	1.000	1.000	0.716	0.888
P32	1.000	0.898	0.811	1.000	1.000	1.000
CF (position)	0.375(12)	0.094(3)	0.438(14)	0.032(1)	0.156(5)	0.094(3)
CF (Amplitude)	1.000	1.000	1.000	1.000	1.000	1.000

Table 6: Calculated Biological Functionalities of P18 and P32.

Sifuvirtide Prototypic Peptide(117-151)	WIEWEREISNYTNQIYEILTESQNNQQRNEKDLLE
---	-------------------------------------

Table 7: Prototypic Peptide of Sifuvirtide.

S/No	Parameter	Enfuvirtide	Sifuvirtide	NHR	CF
1	EIIP	0.815	1.000	0.778	0.281(9)
2	BURA740101	0.467	0.639	0.479	0.562(18)
3	PONP800104	0.745	0.764	1.000	0.283(13)
4	PRAM900102	0.648	0.785	0.821	0.283(13)
5	ARGP820101	0.810	1.000	0.915	0.283(13)
6	ENGD860101	0.637	0.851	0.907	0.283(13)
7	FASG890101	0.841	0.885	1.000	0.283(13)
8	JURD980101	0.828	0.343	0.237	0.283(13)
9	WOLR790101	0.858	0.290	0.208	0.283(13)

Table 8: Calculated Biological Functionalities of Enfuvirtide, Sifuvirtide and NHR.

(amino acids parameters) describe the therapeutic value of the vaccines produced. In both peptides, it is disclosed all interactions by means of the amino acids parameter engaged are above 50%, which appears to suggest that both could be suitably used in the on-going study for the design of anti-Malaria vaccine.

Preliminary Clinical Studies: HIV Fusion Inhibitors

Sifuvirtide, a product of Biomedical engineering is claimed to be more potent, highly effective against Enfuvirtide resistant strains, safer and better tolerated than the Enfuvirtide also called T20 [32]. Prototypic Peptide 7 has been obtained from gp41 of HIV-1 subtype E [37,38]. By means of Biomedical engineering approaches, it was re-design to obtain Sifuvirtide through the introduction of the salt bridge.





Figure 9: CHAM830107-based Common Information Spectrum (CIS) of P18 and P32.

Figure 10: Common Information Spectrum of Enfuvirtide, Sifuvirtide and NHR using PONP800104 and ARGP820101 parameters.




Figure 11: Common Informational Spectrum of Enfuvirtide, Sifuvirtide and NHR using EIIP and BUR740101 parameters.

This led to increased helicity, stability of the Six Helix Bundle (coiled-coil) formed by the anti-HIV/AIDS peptides and the target protein, N-terminus Heptad Repeat (NHR). This determines the anti-HIV/AIDS potency [37,38].

Biomedical engineering design of the Sifuvirtide was achieved by the introduction of the charged amino acids namely, glutamic acid and lysine. Also to provide hydrophobicity pocket, Glutamic acid (E) at position 119 (shown in Table 7 as red, bold and elongated alphabetic code) was replaced with Threonine (T) [32]. Furthermore, Serine was added to the N-terminus so as to increase its stability [32].

As demonstrated above, Biomedical engineering of the Prototypic Peptide 7 into the Sifuvirtide altered the helicity and hydrophobicity properties and as such, three Alpha Helix-related amino acids parameters namely, BUR740101, PONP800104, and PRAM900102 are engaged in this study. Furthermore, five Hydrophobicity-based Amino Acids Parameters including ARGP820101, ENGD860101, FASG890101, JURD980101 and WOLR790101 are employed in the calculation of the pharmacological activity of the two Fusion inhibitors (anti-HIV/AIDS agents). The target protein, N-terminus Heptad Repeat (NHR) is also investigated to show its contribution in interaction. The outcomes of the bio-recognition, bio-affinity and other interactions computationally investigations are as shown in Table 8 using 9 parameters.

Results and Discussions: HIV Fusion Inhibitors

The outcome of the Common Informational Spectrum (CIS) of

the Enfuvirtide, Sifuvirtide and NHR shown in Table 8 using nine Amino Acids Parameters reveals that the CF is at 0.283(13) for all as exemplified by the hydrophobicity-based parameter PONP800104 and Alpha Helix-base ARGP820101 Figure 10 except for the EIIP which is at 0.281(9), and BUR740101 at 0.562(18) though another peak is noticed at position 9 (Figure 11).

From the nine parameters used and shown in Table 8, the amplitude values for the Spectral Characteristics of the Sifuvirtide (using EIIP) is 1.00 (Table 8), which appears to suggest 100% affinity while that of the Enfuvirtide, is 0.815 (Table 8) suggesting 81.5% affinity.

Except for Amino Acids Parameter with descriptor name WOLR79010 (Table 8), Sifuvirtide demonstrated higher amplitude values hence higher biological functionality than the Enfuvirtide. However, based on the nine parameters engaged, the average percentage (%) calculated Biological Functionalities in the Enfuvirtide (72.33%) appear to be less than that obtained in the Sifuvirtide (85.42%) as shown in Table 8. This appears suggest that Sifuvirtide may possess higher therapeutic value as claimed in the preliminary clinical studies. This study is based on the amino acids parameters engaged which are considered to determine the mechanism of actions of the two Fusion Inhibitors of the HIV. Clinically, Sifuvirtide has been suggested to exhibit better efficacy than the Enfuvirtide [32]. Such clinical findings include that the fact that it has six fold higher HIV fusion inhibitory activity. It has been shown that Sifuvirtide not only forms Six Helix bundle (SHB) with target protein (NHR) but blocks other peptides. This is unlike Enfuvirtide.

These claims have preliminarily authenticated by means of CD spectroscopy, a clinical approach. CD spectroscopy has indicated 93% alpha helical content for Sifuvirtide while the Enfuvirtide has none [32]. These factors may have been responsible for the claim that Sifuvirtide is more efficacious as it provides more interaction (average of 85.42%, as shown in Table 8) with the NHR (Table 8), which is known to possess hydrophobic pockets. As noted in Table 8, NHR which interacts with the two anti-HIV/AIDS drugs (Enfuvirtide and Sifuvirtide) offers high level of interaction (87.07%) to these antiretroviral agents.

These findings appear to agree with the calculated Biological functionalities carried out in this study using 9 amino acids parameters that relate to the physiologic indices employed in the clinical experiments (Helicity and Hydrophobicity).

Conclusions

Calculating Biological functionalities by means of Digital Signal Processing techniques appears to be beneficial in the design and development of drugs and vaccines. The introduction of Reverse Vaccinology in the designing of vaccines has resulted in the use of protein fragments or peptides as starter materials for vaccine design. Calculating the Biological characteristics of these peptides rather than obtaining results through clinical experimentation remains a more rational approach. Also, comparisons of Pharmacological activities of the peptide-based drugs by means of calculating their Biological functionalities are a faster and resource saving approach.

Two Digital Signal Processing techniques namely Resonant Recognition Model (RRM) and Informational Spectrum Model (ISM) are employed in calculating the Biological behaviours of two peptides (P18 and P32) which are being investigated for possible incorporation into the materials for the designing of anti-Malaria vaccines. Also, the Biological features of two Fusion inhibitors known as Enfuvirtide (T20) and Sifuvirtide are calculated and the outcomes compared with initial

clinical findings. Furthermore, studying of the effect of mutations on the Biological functionalities by delivering them numerically also quickens the application of the outcomes.

Our results revealed that both P18 and P32 share maximum affinity (100%) which seems to suggest that they offer high specificity that could result in the production of an appropriate antibody. Other interactions studied by means of amino acids parameters engaged which relate potency (sensitivity and neutralization power) of the antibody produced are also found to be high.

Furthermore, our findings disclose that Sifuvirtide (0.8542) has higher average amplitude, which suggest higher interaction (85.42%) than the Enfuvirtide with average amplitude of 0.7233 suggesting 72.33% interaction. This is based on the 9 amino acids parameters which are associated with the physiologic indices clinically, examined indicating higher interaction. This therefore suggests that Sifuvirtide may be more efficacious based on the 9 amino acids parameters studied. This result is found to correlate with clinically derived outcome. However, these results are not interpreted in terms of Pharmacokinetics and Pharmacodynamic activities like solubility, absorption, shelf-life, toxicity, distribution, excretion, etc. As a result, our findings do not suggest acceptability of the Sifuvirtide for the HIV/AIDS management.

This study therefore appears to demonstrate that calculating Biological functionalities is an easier approach to comparing Pharmacological activities of drugs. It also helps determine the Biological activities of peptide components of drugs and vaccines. Manipulation of the amino acids sequences for optimal Biological activities can only be simplified when their Biological functionalities are known and better, delivered or presented in numerical terms.

References

- Smith SW (2002) The Scientist and Engineer's Guide to digital signal processing. California Technical Publishing.
- Cosic I (1994) Macromolecular bioactivity: Is it resonant interaction between macromolecules?-Theory and applications. IEEE Trans Biomed Eng 41: 1101-1114.
- Nwankwo N, Seker H (2010) A signal processing-based bioinformatics approach to assessing drug resistance: human immunodeficiency virus as a case study. Conf Proc IEEE Eng Med Biol Soc 2010: 1836-1839.
- Veljkovic V, Cosic I, Dimitrijevic B, Lalovic D (1985) Is it possible to analyze DNA and protein sequence by the methods of digital signal processing? IEEE Trans Biomed Eng 32: 337-341.
- Veljkovic V, Niman HL, Glisic S, Veljkovic N, Perovic V, et al. (2009) Identification of hemagglutinin structural domain and polymorphisms which may modulate swine H1N1 interactions with human receptor. BMC Struct Biol 9: 62.
- Veljkovic V, Veljkovic N, Muller CP, Muller S, Glisic S, et al. (2009) Characterization of conserved properties of hemagglutinin of H5N1 and human influenza viruses: possible consequences for therapy and infection control. BMC Struct Biol 9: 21.
- Andersen CAF, Brunak S (2004) Representation of protein-sequence information by amino acid subalphabets. AI Magazine 25: 97-104.
- Tomii K, Kanehisa M (1996) Analysis of amino acid indices and mutation matrices for sequence comparison and structure prediction of proteins. Protein Eng 9: 27-36.
- Fernandez L, Caballero J, Abreu JI, Fernandez M (2007) Amino acid sequence autocorrelation vectors and bayesian-regularized genetic neural networks for modeling protein conformational stability: gene V protein mutants. Proteins 67: 834-852.
- Kyte J, Doolittle RF (1982) A simple method for displaying the hydropathic character of a protein. J Mol Biol 157: 105-132.
- Wilce MCJ, Aguilar MI, Hearn MTW (1995) Physicochemical basis of amino acid hydrophobicity scales: Evaluation of four new scales of amino acid hydrophobicity coefficients derived from RP-HPLC of peptides. Anal Chem, 67: 1210-1219.
- Prabhakaran M (1990) The distribution of physical, chemical and conformational properties in signal and nascent peptides. Biochem J 269: 691-696.
- Woods KR, Hopp TP (1981) Prediction of protein antigenic determinants from amino acid sequences. Proc Natl Acad Sci U S A 78: 3824-3828.
- Pique ME, Tainer JA, Getzoff ED, Kuhn LA, Swanson CA (1995) Atomic and residue hydrophilicity in the context of folded protein structures. Proteins 23: 536-547.
- Levitt M, Koehl P (1999) Structure-based conformational preferences of amino acids. Proc Natl Acad Sci U S A 96: 12524-12529.
- Nussinov R, Kumar S, Tsai CJ (2000) Factors enhancing protein thermostability. Protein Eng 13: 179-191.
- Nair AS, Sreenadhan SP (2006) A coding measure scheme employing electron-ion interaction pseudopotential (EIIP). Bioinformation 1: 197-202.
- Frohlich H (1968) Long-range coherence and energy storage in biological systems. Int J Quantum Chem 2: 641-649.
- Veljkovic V, Slavic I (1972) Simple general-model Pseudopotential. Phys Rev Lett 29: 105-107.
- Veljkovic V (1973) The dependence of the fermi energy on the atomic number. Phys Lett A 45: 41-42.
- Doliana R, Veljkovic V (2008) EMILINs interact with anthrax protective antigen and inhibit toxin action in vitro. Matrix Biol 27: 96-106.
- Pirogova E, Akay M, Cosic I (2009) Investigating the interaction between oncogene and tumour suppressor protein. IEEE Trans Inf Technol Biomed 13: 10-15.
- Bajic IV, Bajic VB (1998) Some problems in application of information spectrum method and resonant recognition model for cross-spectral analysis of DNA/RNA sequences. Proc of Communications and Signal Processing 219-224.
- Cooley JW, Tukey JW (1965) An algorithm for the machine calculation of complex fourier series. Math Comput 19: 297-301.
- Aydin N, Haris PI, Chrysostomou C, Seker H (2010) Complex resonant recognition model in analysing influenza a virus subtype protein sequences. IEEE ITAB 1-4.
- Vaidyanathan PP (2004) Genomics and proteomics: A signal processor's tour. Proc of IEEE Circuits and Systems Society 4: 6-29.
- Nwankwo N, Seker H (2010) Assessment of the binding characteristics of human immunodeficiency virus type 1 glycoprotein120 and host cluster of differentiation4 using digital signal processing. IEEE International Conference on BIBE 2010: 289-290.
- Chatterjee S, Wery M, Sharma P, Chauhan VS (1995) A conserved peptide sequence of the Plasmodium falciparum circumsporozoite protein and antipeptide antibodies inhibit Plasmodium berghei sporozoite invasion of Hep-G2 cells and protect immunized mice against P. berghei sporozoite challenge. Infect Immun 63: 4375-4381.
- Cochrane JH (1997) Time Series for Macroeconomics and Finance. Spring.
- Ollivier E, Viari A, Soldano H (1990) A scale-independent signal processing method for sequence analysis. Comput Appl Biosci 6: 71-80.
- Jain E, Bairoch A, Duvaud S, Phan I, Redaschi N, et al. (2009) Infrastructure for the life sciences: design and implementation of the UniProt website. BMC Bioinformatics 10: 136.
- He Y, Xiao Y, Song H, Liang Q, Ju D, et al. (2008) Design and evaluation of sifuvirtide, a novel HIV-1 fusion inhibitor. J Biol Chem 283: 11126-11134.
- Sinnis P, Takacs B, Clavijo P, Santos MJ, Nussenzweig V, et al. (1992) The basolateral domain of the hepatocyte plasma membrane bears receptors for the circumsporozoite protein of Plasmodium falciparum sporozoites. Cell 70: 1021-1033.
- Frevert U, Galinski MR, Hugel F, Allon N, Schreier H, et al. (1998) Malaria circumsporozoite protein inhibits protein synthesis in mammalian cells. EMBO J 17: 3816-3826.
- Pinzon-Ortiz C, Friedman J, Esko J, Sinnis P (2001) The binding of the circumsporozoite protein to cell surface heparan sulfate proteoglycans is required for plasmodium sporozoite attachment to target cells. J Biol Chem 276: 26784-26791.
- Pokarowska M, Kolinski A, Katayama T, Kanehisa M, Kawashima S, et al. (2008) AAindex: amino acid index database. Nucleic Acids Res 36: D202-205.
- Lu M, Kallenbach NR, Spek EJ, Bui AH (1998) Surface salt bridges stabilize the GCN4 leucine zipper. Protein Sci 7: 2431-2437.
- Marqusee S, Baldwin R L (1987) Helix stabilization by glu- lys+ salt bridges in short peptides of de novo design. Proc Natl Acad Sci U S A 84: 8898-8902.